



Service Priorities and Programmes
Electronic Presentations

Convention ID: 139

Submitting author: Dr David CHEUNG

Post title: Resident Specialist, QMH, HKWC

Initiating Insulin Injection for Poorly Controlled DM Patients - Role of Family Physicians at Risk Assessment and Management Program - Associate Consultant (RAMP AC) Clinic of Hong Kong West Cluster

Cheung D, Sze HH, Wang J, Kwong A, Ko W, Tsui WS

Department of Family Medicine and Primary Healthcare

Queen Mary Hospital

Keywords:

Primary Healthcare

DM control

Introduction

Risk Assessment and Management Program - Associate Consultant (RAMP-AC) clinic was established since 2010. It was managed by the Department of Family Medicine and Primary Healthcare of Hong Kong West Cluster (HKWC). Patients with poor DM control in General Out Patient Clinic (GOPC) setting were recruited under the program for insulin initiation. This program involved patients whose:

1. HbA1c \geq 8.5% and categorized as very high risk
2. HbA1c \geq 10% for any categorized risk level
3. Oral anti-diabetic drug (OAD) failure and needs insulin initiation (i.e. HbA1c persistently $>$ 8.5%, on maximum dose)

Objectives

To review the efficacies of insulin initiation of HKWC RAMP AC clinic

Methodology

Statistics of all patients recruited by HKWC RAMP AC clinic from 1/2015 to 12/2015 and starting insulin were reviewed. BP control, Hba1c control and LDL control were the objective markers for the efficacy of service.

Result

From 1/1/2015 to 31/12/2015, there were 318 patients recruited into RAMP AC clinic for insulin initiation. Altogether there were 215 patients on protaphane, 82 patients on mixtard, 12 patients on insulin glargine, 4 patients on insulin aspart and mixtard, 4 patients on actrapid and mixtard, 1 patient on protaphane and mixtard. The mean consultations were 10 times. The mean durations of consultation were 17 months.

When comparing pre- and post- treatment:

The pre- vs post- treatment mean Hba1c = 9.1% vs 8.0%

The pre- vs post- treatment mean LDL = 2.3mmol/L vs 2.1mmol/L

The pre- vs post- treatment mean systolic BP = 130mmHg vs 129mmHg

The pre- vs post- treatment mean diastolic BP = 77mmHg vs 76mmHg

Dramatic improvements were shown at all objective outcomes and further titration of

their insulin doses would be continued at GOPC after discharged from RAMP AC clinic.

From the result we also knew the the time needed to achieve targets for DM patients on insulin were within 1.5 yrs of consultations or 10 visits.

Since there was significant improvement in all objective outcomes after initiating insulin at RAMP AC clinic, it was proved to be an important and effective provider for the service. Family physicians can handle such DM patients at primary care.