



瑪麗醫院
Queen Mary Hospital

尼古丁口含糖補充劑 使用資訊

Nicotine
Replacement
Therapy-
Lozenge
Information
Sheet




無煙新天地
Smoking Counselling & Cessation Centre



Department of Family Medicine & Primary Healthcare (HKWC)
家庭醫學及基層醫療部 (港島西聯網)

使用方法：



注意事項：



應全面停止吸煙



可多飲清水



每天不可多過 20 塊
請勿咬碎口含糖


使用口含糖前後十五分鐘，應避免飲酸性飲料，例如：咖啡、果汁、汽水



若有疑問，可於星期一至五 上午九時至
下午五時致電 查詢：

中西區	
西營盤賽馬會普通科門診診所	2859 8203
中區健康院普通科門診診所	2545 1485
堅尼地城賽馬會普通科門診診所	2817 3215
風險評估及治理診所	2589 2465
護理及專職診所	2589 2490
東華醫院普通科門診診所	2589 8518
南區	
香港仔賽馬會普通科門診診所	2555 0381
鴨脷洲普通科門診診所	2518 5610
風險評估及治理診所	2518 3250
瑪麗醫院專科大樓6樓	22554186

Method to use:



Points to remember:



Should stop smoking completely



Plain fluid intake is advised



Do not use more than 20 pieces in a day
Do not chew the lozenge

Avoid **acidic beverage** such as coffee, soft drink, fruit juice **before and after 15 minutes**, as this will affect the absorption of nicotine.



For any enquiries, please feel free to contact us on Monday to Friday 9am to 5pm


Central & Western District	
Sai Ying Pun Jockey Club GOPC	2859 8203
Central District Health Centre GOPC	2545 1485
Kennedy Town Jockey Club GOPC	2817 3215
Risk Assessment Management Clinic	2589 2465
Nurse & Allied Health Clinic	2589 2490
Tung Wah Hospital GOPC	2589 8518
Southern District	
Aberdeen Jockey Club GOPC	2555 0381
Ap Lei Chau GOPC	2518 5610
Risk Assessment & Management Clinic	2518 3250
Queen Mary Hospital Specialist Clinic S6	22554186

Should you have any queries, please contact your Health Care Provider.

Information provided by Department of Family Medicine and Primary Healthcare, Hong Kong West Cluster

如有任何查詢，請聯絡醫護人員

以上資料由港島西聯網家庭醫學及基層醫療部提供

 瑪麗醫院 Queen Mary Hospital	Pamphlet Topic & Department Code
Ownership	QMH/Family Medicine-7
Last review/revision	May/2023
Approval	HKWC Information Pamphlet Working Group
Distribution	As requested